Name:

Address:

Email:

Provider Number:

My Heart Cardiology Referral Request Form

* Healthlink EDI: myhcardi www.myheartcardiology.com.au ☑ referral@myheartcardiology.com.au

 147 Moreland Road, Coburg 2 38-40 Gap Rd, Sunbury

Suite 8.3, 89 Bridge Rd, Richmond

VIC 3058 VIC 3429

 ▼ Tel: (03) 9854 6444 Fax: (03) 9854 6445 S Neal Street, Gisborne

2 182 Station Rd, New Gisborne

VIC 3437

VIC 3438

VIC 3121

Bartia.	 		

	Putient 3 details							
Clinical Consultation Clinical History: Diabetes Mellitus Hypertension Dyslipidaemia Family history of CVD Obesity Dr Nigel Sutherland Dr Omar Al-Mukhtar Dr Samer Noaman Dr Ahmed Al-Kaisey Next available cardiologist Diagnostic Studies Transthoracic Echocardiogram (TTE) (once every two years) Symptoms or signs of cardiac foilure Ventricular hypertrophy or dysfunction Congenital heart disease Pulmonary hypertension Before starting drugs which require cardiac monitoring Exercise Stress Echocardiogram (once every two years) Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Chest pain or angina Exercise Stress Echocardiogram (once every two years) Exercise Stress Echocardiogram (once every to onalize surgery/intervention on angiography Exercise Stress Echocardiogram (o	Name:	Date of Birth:	Medicare Nur	mber:				
Clinical Consultation Clinical History: Clinical Al-Kaisey: Clone Heart tumour Conditional Heart disease Clinical History: Clone Heart tumour Clone every two years Clone Heart tumour Conditional Heart disease Clinical History: Clone Heart History: C	Phone/Mobile:	Address:						
Clinical History: Diabetes Mellitus Hypertension Oyslipidaemia Family history of CVD Obesity	Clinical Consultation							
Transthoracic Echocardiogram (TTE) (once every two years) Symptoms or signs of cardiac failure Ventricular hypertrophy or dysfunction Volvular, aortic, pericardial, thrombotic or embolic disease Before starting drugs which require cardiac monitoring Exercise Stress Echocardiogram (once every two years) Chest pain or angina Assessment of coronary lesions on angiography Exertional dyspnoea Pre-operative assessment prior to major non-cardiac surgery Suspected silent myocardial ischaemia Pre-operative assessment prior to cardiac surgery/intervention Known coronary disease & evolving symptoms ECG suggestive of coronary disease Cardiac Device Checks Pacemaker Defibrillator Device's company make if known: 12 Lead ECG (trace and report) Heart Rhythm Monitor (please choose duration) 1 Doy 7 Doys Up to 28 Doys Indications: Palpitations Syncope Arrhythmia				□ Diabetes Mellitus□ Hypertension□ Dyslipidaemia□ Family history of CVD				
Transthoracic Echocardiogram (TTE) (once every two years) Symptoms or signs of cardiac failure	☐ Dr Nigel Sutherland ☐ Dr Omar Al-Muk	chtar Dr Samer Noaman	☐ Dr Ahmed Al-Kaisey	Next available cardiologist				
Symptoms or signs of cardiac failure Ventricular hypertrophy or dysfunction Congenital heart disease Valvular, aortic, pericardial, thrombotic or embolic disease Pulmonary hypertension Before starting drugs which require cardiac monitoring Exercise Stress Echocardiogram (once every two years) Chest pain or angina Assessment of coronary lesions on angiography Exertional dyspnoea Pre-operative assessment prior to major non-cardiac surgery Suspected silent myocardial ischaemia Pre-operative assessment prior to cardiac surgery/intervention Known coronary disease & evolving symptoms ECG suggestive of coronary disease Cardiac Device Checks Pacemaker Defibrillator Device's company make if known: 12 Lead ECG (trace and report) Heart Rhythm Monitor (please choose duration) 1 Day 7 Days Up to 28 Days Indications: Palpitations Syncope Arrhythmia	Diagnostic Studies							
□ Cardiac Device Checks □ Pacemaker □ Defibrillator Device's company make if known: □ 12 Lead ECG (trace and report) □ Heart Rhythm Monitor (please choose duration) □ 1 Day □ 7 Days □ Up to 28 Days □ Indications: □ Palpitations □ Syncope □ Arrhythmia □ 24-hour Ambulatory Blood Pressure Monitor (once every 12 months)	Symptoms or signs of cardiac failure Ventricular hypertrophy or dysfunction Valvular, aortic, pericardial, thrombotic or embolic disease Before starting drugs which require cardiac monitoring Exercise Stress Echocardiogram (once every two years) Chest pain or angina Assessment of coronary lesions on angiography Exertional dyspnoea Pre-operative assessment prior to major non-cardiac surgery/intervention							
Pacemaker Defibrillator Device's company make if known: 12 Lead ECG (trace and report) Heart Rhythm Monitor (please choose duration) 1 Day 7 Days Up to 28 Days Indications: Palpitations Syncope Arrhythmia 24-hour Ambulatory Blood Pressure Monitor (once every 12 months)								
Heart Rhythm Monitor (please choose duration) 1 Day 7 Days Up to 28 Days Indications: Palpitations Syncope Arrhythmia 24-hour Ambulatory Blood Pressure Monitor (once every 12 months)	Pacemaker Defibrillator Device's company make if known:							
☐ 1 Day ☐ 7 Days ☐ Up to 28 Days Indications: ☐ Palpitations ☐ Syncope ☐ Arrhythmia ☐ 24-hour Ambulatory Blood Pressure Monitor (once every 12 months)								
·	1 Day 7 Days Up to 28 Days Indications: Palpitations Syncope Arrhythmia							

Phone:

Signature:

Fax:

My Heart Cardiology Referral Request Form

★ Healthlink EDI: myhcardi

www.myheartcardiology.com.au

□ referral@myheartcardiology.com.au

 ▼ Tel: (03) 9854 6444 Fax: (03) 9854 6445 2 38-40 Gap Rd, Sunbury 🙎 5 Neal Street, Gisborne 2 182 Station Rd, New Gisborne

2 147 Moreland Road, Coburg

Suite 8.3, 89 Bridge Rd, Richmond

VIC 3429 VIC 3437 VIC 3438

VIC 3121

VIC 3058

Instructions for Patients

Exercise Stress Echocardiogram & Stress ECG

- A stress echo test measures the performance of your heart under stress. It provides valuable information about blood flow to the heart muscle, heart valve function, and overall cardiac health. During the test, images of your heart are taken before and after exercise on a treadmill (this is for stress echo only).
- The test takes about 20-30 minutes.
- Please wear comfortable exercise clothing and shoes for the test.
- If you take any of the following medications, then you need to stop them for 24-48 hours prior to the test:

NOTE: Consult your doctor before discontinuing any medications.

- Atenolol (Noten)
- Metoprolol (Minax)
- Bisoprolol (Bicor)

- Nebivolol (Nebilet)
- Propranolol (Inderal)
 - Carvedilol (Dilatrend)

- Labetalol
- Sotalol
- Diltiazem (Cardizem) or Verapamil (Isoptin)

Transthoracic Echocardiogram

- Echocardiography is a painless procedure that uses ultrasound technology to produce real-time images of the heart. It allows us to assess the heart's size, shape, and how well it pumps blood.
- The test is typically performed by a specially trained technician and takes about 30-40 minutes.

12 Lead ECG

- This simple test records the electrical signals generated by the heartbeat. These signals are captured by electrodes (small adhesive patches placed on the skin of the chest, arms, and legs).
- This test takes 5 minutes.

Heart Rhythm Monitors

- Holter Monitors

A Holter monitor is a small, wearable device that records your heart's electrical activity for 24 to 48 hours (or longer). It helps us detect irregular heart rhythms that may not appear during a standard ECG in the clinic. While wearing the monitor, you can continue your normal daily activities, but it's essential to note any symptoms such as dizziness, palpitations, or chest discomfort in the provided diary. This information helps us correlate your symptoms with any changes in your heart rhythm.

- HeartBug Monitors

- The HeartBug is a compact, user-friendly device designed to monitor your heart rhythm for extended periods up to several weeks. It's particularly useful for detecting infrequent arrhythmias.
- The HeartBug is simple to apply and comfortable to wear. With this advanced technology, we can gather detailed insights into your heart's activity, ensuring an accurate diagnosis and personalized care plan.

My Heart Cardiology Locations

Richmond Suite 8.3, 89 Bridge Rd, Richmond VIC 3121	Gisborne 5 Neal Street, Gisborne VIC 3437
Coburg 147 Moreland Road, Coburg VIC 3058	New Gisborne 182 Station Rd, New Gisborne VIC 3438
Sunbury 38-40 Gap Rd. Supbury VIC 3429	

referral@myheartcardiology.com.au Healthlink EDI: myhcardi Tel: (03) 9854 6444 Fax: (03) 9854 6445